POLIO TIMELINE





An Egyptian stele portraying Roma the Doorkeeper with a withered leg reveals that poliomyelitis (polio) is an ancient disease



1 7 8 9

British doctor,
Michael Underwood,
attempted the first
known clinical description
of polio, calling it a
"debility of the lower
extremities"






1840

Polio was reported to be contagious by German doctor, Jacob von Heine

Late 'sequelae' (effects)
of polio first mentioned
by French neuropathologist
Jean-Martin Charcot







1894

First polio epidemic in the United States of America

Gould's Illustrated Medical Dictionary defined polio as a "paralysis of certain muscle groups or of an entire limb"






1905

A series of experiments saw Swedish doctor, Ivar Wickman, outline the contagious nature of polio and later categorise the disease into three different types

Austrian doctors,
Karl Landsteiner and
Erwin Popper, discovered
that the infectious agent in
polio was a virus





Polio first became a notifiable disease in Tasmania, with all Australian states and territories effecting this requirement by 1922





1 9 3 2

Australian Sister Kenny initiated a polio treatment that promoted passive and active movements in patients. At the time, this method was deemed controversial as it went against immobilisation recommendations.

However, this treatment is now considered to be the forerunner of modern physiotherapy.

Edward Thomas Both developed Australia's first version of the "Iron Lung"




This negative pressure ventilator enables a person to breathe when normal respiratory muscle control has been lost



North American fundraising organisation "March of Dimes" was established by Franklin Delano Roosevelt to develop a polio vaccine



Australia recorded its highest incidence of paralytic polio (39.1 per 100,000 population)





North America's first mass polio vaccination program was a public health tragedy.

An improperly inactivated vaccine, produced by Cutter Laboratories in California, resulted in 40,000 cases of polio leaving 200 children with varying degrees of paralysis and killing 10.



THE


Cutter Incid 0

HOW AMERICA'S
FIRST POLIO
VACCINE LED TO
THE GROWING
VACCINE CRISIS

Paul Offit, M.D.

Salk inactivated polio vaccine (IPV) introduced in Australia








After working with Dr Salk to develop the polio vaccine, Dr Percival (Val) Bazeley returned to Australia to pioneer its production at the Commonwealth Serum Laboratories (CSL)




Australia's last polio epidemic





Sabin oral polio vaccine (OPV) introduced in Australia







1970°s

Many polio survivors began reporting new problems






1972

Australia's last case of wild polio virus

Australia's Sir Clem Renouf, as President of Rotary International in 1978 and 1979, led the international campaign to vaccinate every child against polio. This resulted in a partnership between the World Health Organisation (WHO), United Nations Children's Fund (UNICEF), US Centers for Disease Control and Prevention (CDC) and Rotary International, and launched the Global Polio Eradication Initiative in 1988.




LATE 1980's

Polio Networks established across Australia



Australia declared polio free by WHO





2007

Polio Flies into Australia

Health authorities launched a nationwide polio alert and spent the next two weeks tracking down passengers on the same flight as a man infected with polio.

The 22-year-old Pakistani man was diagnosed with the infection, the first case reported in Australia in 21 years, after returning to Australia from a holiday in Pakistan.

Polio Australia established at a national conference for State Polio Networks







POLIO AUSTRALIA INCORPORATED

Representing polio survivors throughout Australia



"We're Still Here!"

~ and living in your family tree or memory ~

We are all someone's



We are polio survivors!

Polio Australia is committed to standardising quality information and service provision across Australia for people living with the Late Effects of Polio

> www.polioaustralia.org.au www.stillhere.asn.au

June Middleton, the world's longest surviving polio patient in an iron lung (60 years) died in Melbourne at the age of 83



The number of polio cases worldwide has decreased by more than 99% from 350,000 in 1988 to fewer than 420 cases in 2013. The number of endemic countries has also decreased from over 125 in 1988 to just 3 – Afghanistan, Nigeria and Pakistan. However, reinfected countries include Equatorial Guinea, Iraq, Cameroon, Syria, Ethiopia, Somalia and Kenya.





2014 & BEYOND

In Australia there are an estimated 400,000 polio survivors.

Support services for survivors of polio will be needed for up to 90 years following the very last case of polio contracted by a child anywhere in the world.

The legacy of polio lives on . . .